

## Prostate Core Biopsy for Pathology/Histology Analysis

<b>Test:</b>	Prostate core biopsy (12 part or 6 part) panel.
<b>Submission Criteria:</b>	Needle cores placed immediately into formalin and placed in a sealed biohazard specimen bag.
<b>Storage and Stability:</b>	N/A
<b>Rejection Criteria:</b>	Rejection criteria include but are not limited to those with: <ol style="list-style-type: none"><li>1. Mismatched requisitions</li><li>2. Specimens without TWO patient identifiers</li><li>3. Specimens submitted without approval</li></ol>
<b>Turnaround Time:</b>	2 days
<b>Clinical Significance:</b>	Differential diagnoses include, but are not limited to: High Grade Prostatic Intraepithelial Neoplasia (HGPIN), Atypical small acinar proliferation, Prostatic Adenocarcinoma
<b>Reference Range:</b>	Not detected for all targets
<b>Method:</b>	Panels are comprised of one or more of the following: Hematoxylin and Eosin Stain Immunohistochemical Staining
<b>Platform:</b>	Roche Ventana BenchMark Ultra for IHC staining
<b>CPT Codes:</b>	Biopsy: 55700  Histopathology: 88300, 88302, 88304, 88305, 88307, 88309, 88311, 88213, 88313, 88314, 88319, 88321, 88325, 88342 (CPT codes are assigned by the pathologist when the case is complete)